

Strange duality, mirror symmetry, and the Leech lattice

Wolfgang Ebeling

Dedicated to Terry Wall

Abstract

We give a survey on old and new results concerning Arnold's strange duality. We show that most of the features of this duality continue to hold for the extension of it discovered by C. T. C. Wall and the author. The results include relations to mirror symmetry and the Leech lattice.

Introduction

More than 20 years ago, V. I. Arnold [Ar] discovered a strange duality among the 14 exceptional unimodal hypersurface singularities. A beautiful interpretation of this duality was given by H. Pinkham [P1] and independently by I. V. Dolgachev and V. V. Nikulin [DN, D3]. I. Nakamura related this duality to the Hirzebruch-Zagier duality of cusp singularities [Na1, Na2].

In independent work in early 1982, C. T. C. Wall and the author discovered an extension of this duality embracing on one hand series of bimodal singularities and on the other hand, complete intersection surface singularities in $\mathbb{C}^4$ [EW]. We showed that this duality also corresponds to Hirzebruch-Zagier duality of cusp singularities.

Recent work has aroused new interest in Arnold's strange duality. It was observed by several authors (see [D4] and the references there) that Pinkham's interpretation of Arnold's original strange duality can be considered as part of a two-dimensional analogue of the mirror symmetry of families of Calabi-Yau threefolds. Two years ago, K. Saito [S] discovered a new feature of Arnold's strange duality involving the characteristic polynomials of the monodromy operators of the singularities and he found a connection with the characteristic polynomials of automorphisms of the famous Leech lattice. Only shortly after, M. Kobayashi [Kob] found a duality of the weight systems associated to the 14 exceptional unimodal singularities which corresponds to Arnold's strange duality. He also related it to mirror symmetry.

In this paper we first review these results.

Then we consider our extension of this duality and examine which of the newly discovered features continue to hold. It turns out that with a suitable construction, Pinkham's interpretation can be extended to a larger class of singularities. In this way, one obtains many new examples of mirror symmetric families of K3 surfaces. We also associate characteristic polynomials to the singularities involved in our extension of the duality and show that Saito's duality

continues to hold. Moreover, in this way we can realize further characteristic polynomials of automorphisms of the Leech lattice. The connection with the Leech lattice seems to be rather mysterious. We discuss some facts which might help to understand this connection. We conclude with some open questions.

We thank the referee for his useful comments.

1 Arnold's strange duality

We first discuss Arnold's original strange duality among the 14 exceptional unimodal hypersurface singularities.

We recall Dolgachev's construction [D1, D2] (see also [L1]) of these singularities. Let $b_1 \leq b_2 \leq b_3$ be positive integers such that $\frac{1}{b_1} + \frac{1}{b_2} + \frac{1}{b_3} < 1$. Consider the upper half plane $\mathbb{H} = \{x + iy \in \mathbb{C} | y > 0\}$ with the hyperbolic metric $\frac{1}{y^2}(dx^2 + dy^2)$ and a solid triangle $\Delta \subset \mathbb{H}$ with angles $\frac{\pi}{b_1}, \frac{\pi}{b_2}, \frac{\pi}{b_3}$. Let Σ be the subgroup of the group of isometries of $\mathbb{H}$ generated by the reflections in the edges of Δ , and let Σ_+ be the subgroup of index 2 of orientation preserving isometries. Then $\Sigma_+ \subset \mathrm{PSL}_2(\mathbb{R})$ and Σ_+ acts linearly on the total space $T\mathbb{H}$ of the tangent bundle on $\mathbb{H}$. The inclusion $\mathbb{H} \subset T\mathbb{H}$ as zero section determines an inclusion $\mathbb{H}/\Sigma_+ \subset T\mathbb{H}/\Sigma_+$ of orbit spaces. Collapsing $\mathbb{H}/\Sigma_+$ to a point yields a normal surface singularity (X, x_0) . This singularity is called a *triangle singularity*. The numbers b_1, b_2, b_3 are called the *Dolgachev numbers* $\mathrm{Dol}(X)$ of the singularity. The scalar multiplication in the fibres of the tangent bundle $T\mathbb{H}$ induces a good $\mathbb{C}^*$ -action on X . A resolution of the singularity (X, x_0) can be obtained by the methods of [OW]. A minimal good resolution consists of a rational curve of self-intersection number -1 and three rational curves of self-intersection numbers $-b_1, -b_2$, and $-b_3$ respectively intersecting the exceptional curve transversely.

By [D1], for exactly 14 triples (b_1, b_2, b_3) the singularity (X, x_0) is a hypersurface singularity. Thus it can be given by a function germ $f : (\mathbb{C}^3, 0) \rightarrow (\mathbb{C}, 0)$ where f is weighted homogeneous with weights w_1, w_2, w_3 and degree N . The corresponding weighted homogeneous functions, weights and degrees are indicated in Table 1. It turns out that these singularities are unimodal and one gets in this way exactly the 14 exceptional unimodal hypersurface singularities in Arnold's classification [Ar]. (The equations in Table 1 are obtained by setting the module equal to zero.)

Let (X, x_0) be one of the 14 hypersurface triangle singularities, and denote by X_t and μ its Milnor fibre and Milnor number respectively. We denote by $\langle \cdot, \cdot \rangle$ the intersection form on $H_2(X_t, \mathbb{Z})$ and by $H = (H_2(X_t, \mathbb{Z}), \langle \cdot, \cdot \rangle)$ the Milnor lattice. A. M. Gabrielov [G] has shown that there exists a weakly distinguished basis of vanishing cycles of H with a Coxeter-Dynkin diagram of the form of Fig. 1. The author [E1] has shown that this diagram even corresponds to a distinguished basis of vanishing cycles (cf. also [E5]). (For the notions of a distinguished and weakly distinguished basis of vanishing cycles see e.g. [AGV]). The numbers p_1, p_2, p_3 are called the *Gabrielov numbers* $\mathrm{Gab}(X)$ of the singularity. Here each vertex represents a sphere of self-intersection number -2 , two vertices connected by a single solid edge have intersection number 1, and two vertices connected by a double broken line have intersection number -2 . Using the results of K. Saito (see [E3, Theorem 3.4.3]), one can see that the Gabrielov numbers are uniquely determined by the singularity. We denote by d the discriminant of H , i.e. the

Table 1: The 14 exceptional unimodal singularities

Name	Equation	N	Weights	Dol	Gab	μ	d	Dual
E_{12}	$x^7 + y^3 + z^2$	42	6 14 21	2 3 7	2 3 7	12	1	E_{12}
E_{13}	$x^5 y + y^3 + z^2$	30	4 10 15	2 4 5	2 3 8	13	-2	Z_{11}
E_{14}	$x^8 + y^3 + z^2$	24	3 8 12	3 3 4	2 3 9	14	3	Q_{10}
Z_{11}	$x^5 + xy^3 + z^2$	30	6 8 15	2 3 8	2 4 5	11	-2	E_{13}
Z_{12}	$x^4 y + xy^3 + z^2$	22	4 6 11	2 4 6	2 4 6	12	4	Z_{12}
Z_{13}	$x^6 + xy^3 + z^2$	18	3 5 9	3 3 5	2 4 7	13	-6	Q_{11}
Q_{10}	$x^4 + y^3 + xz^2$	24	6 8 9	2 3 9	3 3 4	10	3	E_{14}
Q_{11}	$x^3 y + y^3 + xz^2$	18	4 6 7	2 4 7	3 3 5	11	-6	Z_{13}
Q_{12}	$x^5 + y^3 + xz^2$	15	3 5 6	3 3 6	3 3 6	12	9	Q_{12}
W_{12}	$x^5 + y^4 + z^2$	20	4 5 10	2 5 5	2 5 5	12	5	W_{12}
W_{13}	$x^4 y + y^4 + z^2$	16	3 4 8	3 4 4	2 5 6	13	-8	S_{11}
S_{11}	$x^4 + y^2 z + xz^2$	16	4 5 6	2 5 6	3 4 4	11	-8	W_{13}
S_{12}	$x^3 y + y^2 z + xz^2$	13	3 4 5	3 4 5	3 4 5	12	13	S_{12}
U_{12}	$x^4 + y^3 + z^3$	12	3 4 4	4 4 4	4 4 4	12	16	U_{12}

determinant of an intersection matrix with respect to a basis of H .

Arnold has now observed: There exists an involution $X \mapsto X^*$ on the set of the 14 exceptional unimodal singularities, such that

$$\text{Dol}(X) = \text{Gab}(X^*), \quad \text{Gab}(X) = \text{Dol}(X^*), \quad N = N^*, \quad \mu + \mu^* = 24.$$

This is called *Arnold's strange duality*. Note that also $d = d^*$.

H. Pinkham [P1] has given the following interpretation of this duality. (This was independently also obtained by I. V. Dolgachev and V. V. Nikulin [DN, D3].) The Milnor fibre X_t can be compactified in a weighted projective space to a surface with three cyclic quotient singularities on the curve at infinity; a minimal resolution of these singularities yields a K3 surface S . Denote by $G(p_1, p_2, p_3)$ the subgraph of the graph of Fig. 1 which is obtained by omitting the vertices with indices $\mu - 1$ and μ . Let $M(p_1, p_2, p_3)$ be the lattice (the free abelian group with an integral quadratic form) determined by the graph $G(p_1, p_2, p_3)$. Then $H = M(p_1, p_2, p_3) \oplus U$, where U is a unimodular hyperbolic plane (the lattice of rank 2 with a basis $\{e, e'\}$ such that $\langle e, e' \rangle = 1$, $\langle e, e \rangle = \langle e', e' \rangle = 0$) and $\oplus$ denotes the orthogonal direct sum. The dual graph of the curve configuration of S at infinity is given by $G(b_1, b_2, b_3)$. The inclusion $X_t \subset S$ induces a primitive embedding $H_2(X_t, \mathbb{Z}) \hookrightarrow H_2(S, \mathbb{Z})$ and the orthogonal complement is just the lattice $M(b_1, b_2, b_3)$. By [Nik] the primitive embedding of $M(p_1, p_2, p_3) \oplus U$ into the unimodular K3 lattice $L := H_2(S, \mathbb{Z})$ is unique up to isomorphism.

In this way, Arnold's strange duality corresponds to a duality of K3 surfaces. This is a two-dimensional analogue of the mirror symmetry between Calabi-Yau threefolds. This has recently been worked out by Dolgachev [D4]. We give an outline of his construction. Let M be an even non-degenerate lattice of signature $(1, t)$. An M -polarized K3 surface is a pair (S, j) where S is a K3 surface and $j : M \hookrightarrow \text{Pic}(S)$ is a primitive lattice embedding. Here $\text{Pic}(S)$ denotes the Picard group of S . An M -polarized K3 surface (S, j) is called *pseudo-ample* if $j(M)$ contains a pseudo-ample divisor class. We assume that M has a unique embedding into the K3 lattice L and the orthogonal complement $M^\perp$ admits an orthogonal splitting $M^\perp = U \oplus \tilde{M}$. (Dolgachev's construction is slightly more




Figure 1: Coxeter-Dynkin diagram of an exceptional unimodal singularity

general.) Then we consider the complete family $\mathcal{F}$ of pseudo-ample M -polarized K3 surfaces and define its *mirror family* $\mathcal{F}^*$ to be any complete family of pseudo-ample $\tilde{M}$ -polarized K3 surfaces. It is shown in [D4] that this is well defined and that there is the following relation between $\mathcal{F}$ and $\mathcal{F}^*$: The dimension of the family $\mathcal{F}$ is equal to the rank of the Picard group of a general member from the mirror family $\mathcal{F}^*$. In particular, this can be applied to $M = M(b_1, b_2, b_3)$ and $\tilde{M} = M(p_1, p_2, p_3)$ for one of the 14 Dolgachev triples (b_1, b_2, b_3) . See [D4] for further results and references.

It was observed by I. Nakamura [Na1, Na2] that Arnold's strange duality corresponds to Hirzebruch-Zagier duality of hyperbolic (alias cusp) singularities. For details see [Na1, Na2, EW].

2 Kobayashi's duality of weight systems

In his paper [Kob], M. Kobayashi has observed a new feature of Arnold's strange duality which we now want to explain.

A quadruple $W = (w_1, w_2, w_3; N)$ of positive integers with $N \in \mathbb{N}w_1 + \mathbb{N}w_2 + \mathbb{N}w_3$ is called a *weight system*. The integers w_i are called the weights and N is called the degree of W . A weight system $W = (w_1, w_2, w_3; N)$ is called *reduced* if $\gcd(w_1, w_2, w_3) = 1$.

Let $W = (w_1, w_2, w_3; N)$ and $W' = (w'_1, w'_2, w'_3; N')$ be two reduced weight systems. An 3×3 -matrix Q whose elements are non-negative integers is called a *weighted magic square* for (W, W') , if

$$(w_1, w_2, w_3)Q = (N, N, N) \quad \text{and} \quad Q \begin{pmatrix} w'_1 \\ w'_2 \\ w'_3 \end{pmatrix} = \begin{pmatrix} N' \\ N' \\ N' \end{pmatrix}.$$

(In the case $w_1 = w_2 = w_3 = w'_1 = w'_2 = w'_3 = 1$, Q is an ordinary magic square.) Q is called *primitive*, if $|\det Q| = N = N'$. We say that the weight systems W and W' are *dual* if there exists a primitive weighted magic square for (W, W') .

Kobayashi now proves:

Theorem 1 (M. Kobayashi) *Let $W = (w_1, w_2, w_3; N)$ be the weight system of one of the 14 exceptional unimodal singularities. Then there exists a unique (up to permutation) dual weight system W^* . The weight system W^* belongs to the dual singularity in the sense of Arnold.*

Moreover, Kobayashi shows that there is a relation between this duality of weight systems and the polar duality between certain polytopes associated to the weight systems. Such a polar duality was considered by V. Batyrev [Ba] in connection with the mirror symmetry of Calabi-Yau hypersurfaces in toric varieties. We refer to [E6] for a more precise discussion of this relation.

3 Saito's duality of characteristic polynomials

Let $f : (\mathbb{C}^3, 0) \rightarrow (\mathbb{C}, 0)$ be a germ of an analytic function defining an isolated hypersurface singularity (X, x_0) . A characteristic homeomorphism of the Milnor fibration of f induces an automorphism $c : H_2(X_t, \mathbb{Z}) \rightarrow H_2(X_t, \mathbb{Z})$ called the (*classical*) *monodromy operator* of (X, x_0) . It is a well known theorem (see e.g. [Br]) that the eigenvalues of c are roots of unity. This means that the characteristic polynomial $\phi(\lambda) = \det(\lambda I - c)$ of c is a monic polynomial the roots of which are roots of unity. Such a polynomial can be written in the form

$$\phi(\lambda) = \prod_{m \geq 0} (\lambda^m - 1)^{\chi_m} \quad \text{for } \chi_m \in \mathbb{Z},$$

where all but finitely many of the integers χ_m are equal to zero. We note some useful formulae.

Proposition 1 (i) $\mu = \deg \phi = \sum_{m > 0} m \chi_m$.

(ii) If $\sum_{m > 0} \chi_m = 0$ then

$$\phi(1) = \prod_{m > 0} m^{\chi_m}.$$

(iii) $\phi(1) = (-1)^\mu d$.

(iv) $\text{tr } c^k = \sum_{m|k} m \chi_m$.
In particular $\text{tr } c = \chi_1$.

Proof. (i) is obvious. For the proof of (ii) we use the identity

$$(\lambda^m - 1) = (\lambda - 1)(\lambda^{m-1} + \dots + \lambda + 1).$$

To prove (iii), let A be the intersection matrix with respect to a distinguished basis $\{\delta_1, \dots, \delta_\mu\}$ of vanishing cycles. Write A in the form $A = V + V^t$ where V is an upper triangular matrix with -1 on the diagonal. Let C be the matrix

of c with respect to $\{\delta_1, \dots, \delta_\mu\}$. Then $C = -V^{-1}V^t$ (see e.g. [E3, Proposition 1.6.3]). Therefore

$$\phi(1) = \det(I - C) = \det V^{-1}(V + V^t) = (-1)^\mu d.$$

Finally, (iv) is obtained as in [AC2] using the identity

$$\det(I - tc) = \exp(\operatorname{tr}(\log(I - tc))) = \exp\left(-\sum_{k \geq 1} \frac{t^k}{k} \operatorname{tr} c^k\right).$$

This proves Proposition 1.

By A'Campo's theorem [AC1]

$$\operatorname{tr} c = -1.$$

We assume that c has finite order h . This is e.g. true if f is a weighted homogeneous polynomial of degree N . In this case $h = N$. Then $\chi_m = 0$ for all m which do not divide h . K. Saito [S] defines a *dual polynomial* $\phi^*(\lambda)$ to $\phi(\lambda)$:

$$\phi^*(\lambda) = \prod_{k|h} (\lambda^k - 1)^{-\chi_{h/k}}.$$

He obtains the following result.

Theorem 2 (K. Saito) *If $\phi(\lambda)$ is the characteristic polynomial of the monodromy of an exceptional unimodal singularity X , then $\phi^*(\lambda)$ is the corresponding polynomial of the dual singularity X^* .*

For $\phi(\lambda) = \prod_{m|h} (\lambda^m - 1)^{\chi_m}$ we use the symbolic notation

$$\pi := \prod_{m|h} m^{\chi_m}.$$

In the theory of finite groups, this symbol is known as a *Frame shape* [F, CN]. For, if one has a rational finite-dimensional representation of a finite group, then the zeros of the characteristic polynomials of each element of the group are also roots of unity. For a given rational representation, one can thus assign to each conjugacy class of the group its Frame shape. The number

$$\deg(\pi) = \sum m\chi_m$$

is called the *degree* of the Frame shape π .

Let us denote the Frame shape of the dual polynomial $\phi^*(\lambda)$ by π^* . The Frame shapes of the monodromy operators of the 14 exceptional unimodal singularities are listed in Table 2.

To two Frame shapes $\pi = \prod m^{\chi_m}$ and $\pi' = \prod m^{\chi'_m}$ of degree n and n' respectively one can associate a Frame shape $\pi\pi'$ of degree $n+n'$ by concatenation

$$\pi\pi' := \prod m^{\chi_m} \prod m^{\chi'_m} = \prod m^{\chi_m + \chi'_m}.$$

If π and π' are the Frame shapes of the operators $c : H \rightarrow H$ and $c' : H' \rightarrow H'$ respectively, $\pi\pi'$ is the Frame shape of the operator $c \oplus c' : H \oplus H' \rightarrow H \oplus H'$.

Table 2: Frame shapes of the 14 exceptional unimodal singularities

Name	π	π^*	Dual
E_{12}	$2 \cdot 3 \cdot 7 \cdot 42/1 \cdot 6 \cdot 14 \cdot 21$	$2 \cdot 3 \cdot 7 \cdot 42/1 \cdot 6 \cdot 14 \cdot 21$	E_{12}
E_{13}	$2 \cdot 3 \cdot 30/1 \cdot 6 \cdot 15$	$2 \cdot 5 \cdot 30/1 \cdot 10 \cdot 15$	Z_{11}
E_{14}	$2 \cdot 3 \cdot 24/1 \cdot 6 \cdot 8$	$3 \cdot 4 \cdot 24/1 \cdot 8 \cdot 12$	Q_{10}
Z_{12}	$2 \cdot 22/1 \cdot 11$	$2 \cdot 22/1 \cdot 11$	Z_{12}
Z_{13}	$2 \cdot 18/1 \cdot 6$	$3 \cdot 18/1 \cdot 9$	Q_{11}
Q_{12}	$3 \cdot 15/1 \cdot 5$	$3 \cdot 15/1 \cdot 5$	Q_{12}
W_{12}	$2 \cdot 5 \cdot 20/1 \cdot 4 \cdot 10$	$2 \cdot 5 \cdot 20/1 \cdot 4 \cdot 10$	W_{12}
W_{13}	$2 \cdot 16/1 \cdot 4$	$4 \cdot 16/1 \cdot 8$	S_{11}
S_{12}	$13/1$	$13/1$	S_{12}
U_{12}	$4 \cdot 12/1 \cdot 3$	$4 \cdot 12/1 \cdot 3$	U_{12}

In the appendix of [S], Saito notes the following observation: If π is the Frame shape of the monodromy operator of an exceptional unimodal singularity, then the symbol $\pi\pi^*$ of degree 24 is a Frame shape of a conjugacy class of the automorphism group of the Leech lattice. The Leech lattice is a 24-dimensional even unimodular positive definite lattice which contains no roots (see e.g. [E4]). It was discovered by J. Leech in connection with the search for densest sphere packings. Its automorphism group G is a group of order $2^{22}3^95^47^211 \cdot 13 \cdot 23$. The quotient group $\text{Co}_1 := G/\{\pm 1\}$ is a famous sporadic simple group discovered by J. Conway. The Frame shapes of the 164 conjugacy classes of G have been listed by T. Kondo [Kon].

4 An extension of Arnold's strange duality

C. T. C. Wall and the author [EW] have found an extension of Arnold's strange duality. In order to consider this extension, we have to enlarge the class of singularities which we want to discuss.

On the one hand, instead of restricting to the hypersurface case, we can also look at isolated complete intersection singularities (abbreviated ICIS in the sequel). Pinkham has shown [P2] that for exactly 8 triples (b_1, b_2, b_3) the triangle singularities with these Dolgachev numbers are ICIS, but not hypersurface singularities. They are given by germs of analytic mappings $(g, f) : (\mathbb{C}^4, 0) \rightarrow (\mathbb{C}^2, 0)$. They are $\mathcal{K}$ -unimodal singularities and appear in Wall's classification [Wa]. For certain values of the module, the equations are again weighted homogeneous. The corresponding 8 triples (b_1, b_2, b_3) , Wall's names and weighted homogeneous equations with weights (w_1, w_2, w_3, w_4) and degrees (N_1, N_2) are indicated in Table 3.

By [H1], the notion of Milnor fibre can also be extended to ICIS. We assume that (g, f) are generically chosen such that $(X', 0) = (g^{-1}(0), 0)$ is an isolated hypersurface singularity of minimal Milnor number μ_1 among such choices of g . Then the *monodromy operator* of $(X, 0)$ is defined to be the monodromy operator of the function germ $f : (X', 0) \rightarrow (\mathbb{C}, 0)$. By [E3] there exists a distinguished set of generators consisting of $\nu := \mu + \mu_1$ vanishing cycles, where μ is the rank of the second homology group of the Milnor fibre. Again the monodromy operator is the Coxeter element of this set, i.e. the product of the ν reflections corresponding

Table 3: The 8 triangle ICIS

Name	Equations	N	Weights	Dol	Gab	μ	d	Dual
J'_9	$xw + y^2$ $x^3 + yw + z^2$	16 18	6 8 9 10	2 3 10	2 2 2 3	9	-4	$J_{3,0}$
J'_{10}	$xw + y^2$ $x^2y + yw + z^2$	12 14	4 6 7 8	2 4 8	2 2 2 4	10	8	$Z_{1,0}$
J'_{11}	$xw + y^2$ $x^4 + yw + z^2$	10 12	3 5 6 7	3 3 7	2 2 2 5	11	-12	$Q_{2,0}$
K'_{10}	$xw + y^2$ $x^3 + z^2 + w^2$	10 12	4 5 6 6	2 6 6	2 3 2 3	10	12	$W_{1,0}$
K'_{11}	$xw + y^2$ $x^2y + z^2 + w^2$	8 10	3 4 5 5	3 5 5	2 3 2 4	11	-20	$S_{1,0}$
L_{10}	$xw + yz$ $x^3 + yw + z^2$	11 12	4 5 6 7	2 5 7	2 2 3 3	10	11	$W_{1,0}$
L_{11}	$xw + yz$ $x^2y + yw + z^2$	9 10	3 4 5 6	3 4 6	2 2 3 4	11	-18	$S_{1,0}$
M_{11}	$2xw + y^2 + z^2$ $x^3 + 2yw$	8 9	3 4 4 5	4 4 5	2 3 3 3	11	-24	$U_{1,0}$

to the vanishing cycles of the distinguished set of generators. For the 8 triangle ICIS, a Coxeter-Dynkin diagram corresponding to such a distinguished set is depicted in Fig. 2 (cf. [E3]). Let us call the characteristic numbers p_1, p_2, p_3, p_4 of these graphs the *Gabrielov numbers* $\text{Gab}(X)$ of the singularity. They are also indicated in Table 3. Again, using [E3, Theorem 3.4.3] one can see that these numbers are uniquely defined.

On the other hand, instead of starting with a hyperbolic triangle, we can start with a hyperbolic quadrilateral. Let b_1, b_2, b_3, b_4 be positive integers such that

$$\frac{1}{b_1} + \frac{1}{b_2} + \frac{1}{b_3} + \frac{1}{b_4} < 2.$$

One can perform the same construction as above with a solid quadrilateral with angles $\frac{\pi}{b_1}, \frac{\pi}{b_2}, \frac{\pi}{b_3}, \frac{\pi}{b_4}$ instead of a triangle. The resulting normal surface singularities are called *quadrilateral singularities* (with *Dolgachev numbers* b_1, b_2, b_3, b_4) (cf. [L2]). Again these singularities admit a natural good $\mathbb{C}^*$ -action. A minimal good resolution consists of a rational curve of self-intersection number -2 together with four rational curves of self-intersection numbers $-b_1, -b_2, -b_3$, and $-b_4$ respectively intersecting the first curve transversely.

For 6 quadruples (b_1, b_2, b_3, b_4) these singularities are isolated hypersurface singularities. They are bimodal in the sense of Arnold. They can again be defined by weighted homogeneous equations. By [EW] they have a distinguished basis with a Coxeter-Dynkin diagram of the following form: It is obtained by adding a new vertex with number 0 to the graph of Fig. 1 in one of the following two ways:

- (1) It is connected to the vertex μ and to the vertex $p_1 + p_2$ by ordinary lines. We refer to this diagram by the symbol $(p_1, p_2, \underline{p_3})$.
- (2) It is connected to the vertex μ and to the vertices p_1 and $p_1 + p_2 - 1$ by ordinary lines. We refer to this diagram by the symbol $(p_1, \underline{p_2}, \underline{p_3})$.




Figure 2: Coxeter-Dynkin diagram of a triangle ICIS

Table 4: The 6 quadrilateral hypersurface singularities

Name	Equation	N	Weights	Dol	Gab	μ	d	Dual
$J_{3,0}$	$x^9 + y^3 + z^2$	18	2 6 9	2 2 2 3	2 3 <u>10</u>	16	4	J'_9
$Z_{1,0}$	$x^7 + xy^3 + z^2$	14	2 4 7	2 2 2 4	2 4 <u>8</u>	15	-8	J'_{10}
$Q_{2,0}$	$x^6 + y^3 + xz^2$	12	2 4 5	2 2 2 5	3 3 <u>7</u>	14	12	J'_{11}
$W_{1,0}$	$x^6 + y^4 + z^2$	12	2 3 6	2 2 3 3	$\begin{smallmatrix} 2 & 5 & \underline{7} \\ 2 & \underline{6} & \underline{6} \end{smallmatrix}$	15	-12	$\begin{smallmatrix} K'_{10} \\ L_{10} \end{smallmatrix}$
$S_{1,0}$	$x^5 + y^2z + xz^2$	10	2 3 4	2 2 3 4	$\begin{smallmatrix} 3 & 4 & \underline{6} \\ 3 & \underline{5} & \underline{5} \end{smallmatrix}$	14	20	$\begin{smallmatrix} K'_{11} \\ L_{11} \end{smallmatrix}$
$U_{1,0}$	$x^3y + y^3 + z^3$	9	2 3 3	2 3 3 3	4 <u>4</u> <u>5</u>	14	27	M_{11}

The corresponding data are indicated in Table 4. The numbers p_1, p_2, p_3 defined by this procedure are unique except for the singularities $W_{1,0}$ and $S_{1,0}$, where we have two triples each. This follows from [E2, Supplement to Theorem 4.1]. One can also obtain an easier proof by considering the discriminants (and in one case the discriminant quadratic form) of the singularities and the possible determinants of the graphs (and in one case the discriminant quadratic form determined by the graph). The absolute values of the determinants in the respective cases are

- (1) $|d| = 4(p_1p_2 - p_1 - p_2)$,
- (2) $|d| = (p_1 - 1)(p_2 + p_3)$.

For another 5 quadruples (b_1, b_2, b_3, b_4) the quadrilateral singularities with these Dolgachev numbers are ICIS. They are also $\mathcal{K}$ -unimodal and appear in Wall's lists [Wa]. They can also be given by weighted homogeneous equations. These equations and Wall's names are listed in Table 5.

Table 5: The 5 quadrilateral ICIS

Name	Equations	Restrictions	N	Weights
$J'_{2,0}$	$xw + y^2$ $ax^5 + xy^2 + yw + z^2$	$a \neq 0, -\frac{4}{27}$	$\begin{smallmatrix} 8 \\ 10 \end{smallmatrix}$	2 4 5 6
$L_{1,0}$	$xw + yz$ $ax^4 + xy^2 + yw + z^2$	$a \neq 0, -1$	$\begin{smallmatrix} 7 \\ 8 \end{smallmatrix}$	2 3 4 5
$K'_{1,0}$	$xw + y^2$ $ax^4 + xy^2 + z^2 + w^2$	$a \neq 0, \frac{1}{4}$	$\begin{smallmatrix} 6 \\ 8 \end{smallmatrix}$	2 3 4 4
$M_{1,0}$	$2xw + y^2 - z^2$ $x^2y + ax^2z + 2yw$	$a \neq \pm 1$	$\begin{smallmatrix} 6 \\ 7 \end{smallmatrix}$	2 3 3 4
$I_{1,0}$	$x^3 + w(y - z)$ $ax^3 + y(z - w)$	$a \neq 0, 1$	$\begin{smallmatrix} 6 \\ 6 \end{smallmatrix}$	2 3 3 3

Name	Dol	Gab	μ	d	Dual
$J'_{2,0}$	2 2 2 6	2 2 2 <u>6</u>	13	-16	$J'_{2,0}$
$L_{1,0}$	2 2 3 5	$\begin{smallmatrix} 2 & 2 & 3 & \underline{5} \\ 2 & 2 & \underline{4} & \underline{4} \end{smallmatrix}$	13	-28	$\begin{smallmatrix} L_{1,0} \\ K'_{1,0} \end{smallmatrix}$
$K'_{1,0}$	2 2 4 4	$\begin{smallmatrix} 2 & 3 & 2 & \underline{5} \\ 2 & \underline{4} & 2 & \underline{4} \end{smallmatrix}$	13	-32	$\begin{smallmatrix} L_{1,0} \\ K'_{1,0} \end{smallmatrix}$
$M_{1,0}$	2 3 3 4	$\begin{smallmatrix} 2 & 3 & \underline{3} & \underline{4} \\ 2 & \underline{3} & 3 & \underline{4} \end{smallmatrix}$	13	-42	$M_{1,0}$
$I_{1,0}$	3 3 3 3	$\underline{3} \underline{3} \underline{3} \underline{3}$	13	-54	$I_{1,0}$

The singularities $J'_{2,0}$, $L_{1,0}$, $K'_{1,0}$, and $M_{1,0}$ can be given by equations (g, f) where g has Milnor number $\mu_1 = 1$. Coxeter-Dynkin diagrams of these singularities are computed in [E3]. Using transformations as in the proof of [E3, Proposition 3.6.1], these graphs can be transformed to the following graphs. A Coxeter-Dynkin diagram corresponding to a distinguished set of generators is obtained by adding a new vertex to the graph of Fig. 2. It gets the number $p_1 + p_2 + 2$ and the indices of the old vertices with numbers $p_1 + p_2 + 2, p_1 + p_2 + 3, \dots, \nu$ are shifted by 1. New edges are introduced in one of the following ways:

- (1) The new vertex is connected to the vertex $p_1 + p_2 + 1$ and to the vertex with new index $p_1 + p_2 + p_3 + 3$ (old index $p_1 + p_2 + p_3 + 2$) by ordinary lines. We refer to this diagram by the symbol $(p_1, p_2, p_3, \underline{p_4})$.
- (2) The new vertex is connected to the vertex $p_1 + p_2 + 1$ and to the vertices with new indices $p_1 + p_2 + 3$ and $p_1 + p_2 + p_3 + 2$ (old indices $p_1 + p_2 + 2$ and $p_1 + p_2 + p_3 + 1$ respectively) by ordinary lines. We refer to this diagram by the symbol $(p_1, p_2, \underline{p_3}, \underline{p_4})$.
- (3) The new vertex is connected to the vertex $p_1 + p_2 + 1$, to the vertex p_1 , and to the vertex with new index $p_1 + p_2 + p_3 + 2$ (old index $p_1 + p_2 + p_3 + 1$) by ordinary lines. We refer to this diagram by the symbol $(p_1, \underline{p_2}, p_3, \underline{p_4})$.

The absolute values of the determinants of the respective graphs are

- (1) $|d| = 4(p_1 p_2 p_3 - p_1 - p_3)$,
- (2) $|d| = p_1 p_2 (p_3 + p_4 + 2) - p_1 - p_2 - p_3 - p_4$,
- (3) $|d| = p_1 p_3 (p_2 + p_4)$.

Comparing the values of these determinants with the discriminants of the above 4 quadrilateral ICIS, we find that the graphs listed in Table 5 are the only possible graphs of the types (1), (2), or (3) for these singularities. Again, in two cases the numbers p_1, p_2, p_3, p_4 are not uniquely defined.

For the remaining singularity $I_{1,0}$, $\mu_1 = 2$. This singularity can be given by the following equations:

$$\begin{aligned} g(z) &= z_1^2 + z_2^2 + z_3^2 + z_4^3, \\ f(z) &= a_1 z_1^2 + a_2 z_2^2 + a_3 z_3^2 + a_4 z_4^3, \end{aligned}$$

where $a_i \in \mathbb{R}$, $a_1 < a_2 < a_3 < a_4$. For such a singularity H. Hamm [H2] has given a basis of the complexified Milnor lattice $H_{\mathbb{C}} = H \otimes \mathbb{C}$. As in [E3, Sect. 2.3], one can show that the cycles he constructs are in fact vanishing cycles and that there exists a distinguished set $\{\delta_1, \dots, \delta_\nu\}$ of generators for this singularity with the following intersection numbers:

$$\begin{aligned} \langle \delta_{i+1}, \delta_{i+2} \rangle &= -1, \quad i = 0, 2, 4, 6, 8, 10, \\ \langle \delta_{i+1}, \delta_{i+3} \rangle &= \langle \delta_{i+1}, \delta_{i+4} \rangle = 0, \quad \langle \delta_{i+2}, \delta_{i+3} \rangle = \langle \delta_{i+2}, \delta_{i+4} \rangle = 0, \quad i = 0, 4, 8, \\ \langle \delta_{i+1}, \delta_j \rangle &= \langle \delta_{i+3}, \delta_j \rangle = -1, \quad i = 0, 4, 8, \quad 1 \leq j \leq 12, j \neq i+1, i+2, i+3, i+4, \\ \langle \delta_{i+2}, \delta_{2k} \rangle &= \langle \delta_{i+4}, \delta_{2k} \rangle = -1, \quad i = 0, 4, 8, \quad 1 \leq k \leq 6, 2k \neq i+2, i+4, \\ \langle \delta_{i+2}, \delta_{2k-1} \rangle &= \langle \delta_{i+4}, \delta_{2k-1} \rangle = 0, \quad i = 0, 4, 8, \quad 1 \leq k \leq 6, 2k-1 \neq i+1, i+3, \\ \langle \delta_{2k-1}, \delta_{13} \rangle &= -1, \quad \langle \delta_{2k-1}, \delta_{15} \rangle = 0, \quad \langle \delta_{2k}, \delta_{13} \rangle = 0, \quad \langle \delta_{2k}, \delta_{15} \rangle = -1, \quad 1 \leq k \leq 6, \\ \langle \delta_i, \delta_{14} \rangle &= -1, \quad 1 \leq i \leq 12, \\ \langle \delta_{13}, \delta_{14} \rangle &= 0, \quad \langle \delta_{13}, \delta_{15} \rangle = 0, \quad \langle \delta_{14}, \delta_{15} \rangle = 0. \end{aligned}$$

By the following sequence of braid group transformations (for the notation see e.g. [E3]) the distinguished set $\{\delta_1, \dots, \delta_\nu\}$ can be transformed to a distinguished set $\{\delta'_1, \dots, \delta'_\nu\}$ where the Coxeter-Dynkin diagram corresponding to the subset $\{\delta'_1, \dots, \delta'_{12}\}$ is the graph of Fig. 3:

$$\begin{aligned} &\beta_5, \beta_4, \beta_3, \beta_2; \alpha_6, \alpha_7, \alpha_8, \alpha_9, \alpha_{10}, \alpha_{11}; \alpha_5, \alpha_6; \alpha_3, \alpha_4, \alpha_5; \\ &\beta_8, \beta_7, \beta_6; \beta_{10}, \beta_9, \beta_8, \beta_7; \kappa_4, \kappa_5, \kappa_6, \kappa_7, \kappa_8, \kappa_9. \end{aligned}$$

We refer to the Coxeter-Dynkin diagram corresponding to $\{\delta'_1, \dots, \delta'_\nu\}$ by the symbol $(\underline{3}, \underline{3}, \underline{3}, \underline{3})$. This notation is motivated by Fig. 3. We admit, however, that it is somewhat arbitrary.

The corresponding symbols are indicated in Table 5.

If one compares the Dolgachev and Gabrielov numbers of Tables 3 and 4 and of Table 5, then one observes a correspondence between the 8 triangle ICIS and the 6 quadrilateral hypersurface singularities and between the 5 quadrilateral ICIS. The corresponding "dual" singularities are indicated in the last column of each table. Note that this correspondence is not always a duality in the strict sense. For the quadrilateral hypersurface singularity $W_{1,0}$ we have two corresponding ICIS K'_{10} and L_{10} and for $S_{1,0}$ we have the corresponding ICIS K'_{11} and L_{11} . The quadrilateral ICIS $L_{1,0}$ and $K'_{1,0}$ are both self-dual and dual to each other. In the other cases the correspondence is one-to-one. Pinkham also defined "Gabrielov numbers" (in a weaker sense) for the triangle ICIS [P2]




Figure 3: Subgraph of a Coxeter-Dynkin diagram of the singularity $I_{1,0}$

and he already made part of this observation (unpublished). This duality also corresponds to the Hirzebruch-Zagier duality of cusp singularities (see [Na2, EW]).

If one now compares the Milnor numbers of dual singularities, one finds

- for the triangle ICIS versus quadrilateral hypersurface singularities: $\mu + \mu^* = 25$.
- for the quadrilateral ICIS: $\mu + \mu^* = 26$.

(Note that also d and d^* do not coincide in each case.) So one still has to alter something. There are two alternatives:

- (1) subtract 1 for quadrilateral.
- (2) subtract 1 for ICIS.

In [EW] we considered the first alternative. The quadrilateral singularities are first elements ($l = 0$) of series of singularities indexed by a non-negative integer l . We showed that to each such series one can associate a virtual element $l = -1$. We defined for these Milnor lattices, Coxeter-Dynkin diagrams, and monodromy operators, and showed that all features of Arnold's strange duality including Pinkham's interpretation continue to hold. For more details see below.

A new discovery is that the second alternative works as well, and this also leads to an extension of Saito's duality. Recall that the triangle or quadrilateral ICIS with $\mu_1 = 1$ have a Coxeter-Dynkin diagram D which is either the graph of Fig. 2 or an extension of it. By similar transformations as in the proof of [E3, Proposition 3.6.2], this graph can be transformed to a graph containing the subgraph D^b depicted in Fig. 4. (Unfortunately, this proof has to be modified slightly. The correct sequence of transformations is

$$\beta_{\rho-1}, \beta_{\rho-2}, \beta_{\rho-3}, \beta_{\rho-3}, \beta_{\rho-2}, \kappa_{\rho-1}$$

and one has to consider the vertices $\lambda'_{\rho-2}$ and λ'_ρ instead of $\lambda'_{\rho-1}$ and λ'_ρ .) By [E3, Remark 3.6.5], the passage from D to D^b can be considered as a kind of "desuspension". For the singularity $I_{1,0}$, let D^b be the graph of Fig. 3. In each




Figure 4: Reduced Coxeter-Dynkin diagram D^b

case, the new Coxeter-Dynkin diagram D^b has $\mu - 1$ instead of $\mu + \mu_1$ vertices. Denote the corresponding Coxeter element (product of reflections corresponding to the vanishing cycles) by c^b . For the ICIS with $\mu_1 = 1$ one can compute (see [E3, Proposition 3.6.2]) that

$$\pi(c^b) = \pi(c)/1.$$

This means that c^b has the same eigenvalues as c but the multiplicities of the eigenvalue 1 differ by 1. For the singularity $I_{1,0}$ one has $\pi(c) = 6^3/1 \cdot 2^2$ (cf. [H2]), whereas $\pi(c^b) = 3^2 6^2/1^2 2^2$. Note that in any case

$$\text{tr } c^b = -2.$$

The passage from c to c^b corresponds to the passage from the Milnor lattice H of rank μ to a sublattice H^b of rank $\mu^b = \mu - 1$. The corresponding discriminants and discriminant quadratic forms of the lattices H^b are listed in Table 6. Here we use the notation of [EW].

The Frame shapes of the corresponding operators c^b are listed in Table 7.

It turns out that the substitution

$$\mu \mapsto \mu^b, \quad c \mapsto c^b, \quad H \mapsto H^b$$

for the ICIS yields

$$\mu + \mu^* = 24, \quad d = d^*, \quad \pi_{X^*} = \pi_X^*.$$

Moreover, the lattice H admits an embedding into the even unimodular lattice

$$K_{24} = (-E_8) \oplus (-E_8) \oplus U \oplus U \oplus U \oplus U$$

Table 6: Discriminants and discriminant quadratic forms of the lattices H^b

Name	μ^b	d^b	$(H^b)^*/H^b$	dual form	dual
J'_9	8	4	q_{D_4}	q_{D_4}	$J_{3,0}$
J'_{10}	9	-8	$q_{D_4} + q_{A_1}$	$q_{D_4} + w_{2,1}^1$	$Z_{1,0}$
J'_{11}	10	12	$q_{D_4} + q_{A_2}$	$q_{D_4} + w_{3,1}^1$	$Q_{2,0}$
K'_{10}	9	-12	$w_{2,2}^1 + w_{3,1}^{-1}$	$w_{2,2}^{-1} + w_{3,1}^1$	$W_{1,0}$
L_{10}	9	-12	$w_{2,2}^1 + w_{3,1}^{-1}$	$w_{2,2}^{-1} + w_{3,1}^1$	$W_{1,0}$
K'_{11}	10	20	$w_{2,1}^1 + w_{2,1}^1 + w_{5,1}^{-1}$	$w_{2,1}^{-1} + w_{2,1}^{-1} + w_{5,1}^{-1}$	$S_{1,0}$
L_{11}	10	20	$w_{2,1}^1 + w_{2,1}^1 + w_{5,1}^{-1}$	$w_{2,1}^{-1} + w_{2,1}^{-1} + w_{5,1}^{-1}$	$S_{1,0}$
M_{11}	10	27	$w_{3,2}^{-1} + w_{3,1}^{-1}$	$w_{3,2}^1 + w_{3,1}^1$	$U_{1,0}$
$J'_{2,0}$	12	16	$v_1 + v_1$	$v_1 + v_1$	$J'_{2,0}$
$L_{1,0}$ $K'_{1,0}$	12	32	$w_{2,2}^1 + w_{2,3}^{-1}$	$w_{2,2}^1 + w_{2,3}^{-1}$	$L_{1,0}$ $K'_{1,0}$
$M_{1,0}$	12	49	$w_{7,1}^1 + w_{7,1}^{-1}$	$w_{7,1}^1 + w_{7,1}^{-1}$	$M_{1,0}$
$I_{1,0}$	12	81	$w_{3,1}^1 + w_{3,1}^1 + w_{3,1}^{-1} + w_{3,1}^{-1}$	$w_{3,1}^{-1} + w_{3,1}^{-1} + w_{3,1}^1 + w_{3,1}^1$	$I_{1,0}$

Table 7: Frame shapes of the triangle ICIS and quadrilateral singularities

Name	π	π^*	Dual
$J_{3,0}$	$2 \cdot 3 \cdot 18^2/1 \cdot 6 \cdot 9^2$	$2^2 3 \cdot 18/1^2 6 \cdot 9$	J'_9
$Z_{1,0}$	$2 \cdot 14^2/1 \cdot 7^2$	$2^2 14/1^2 7$	J'_{10}
$Q_{2,0}$	$3 \cdot 12^2/1 \cdot 6^2$	$2^2 12/1^2 4$	J'_{11}
$W_{1,0}$	$2 \cdot 12^2/1 \cdot 4 \cdot 6$	$2 \cdot 3 \cdot 12/1^2 6$	K'_{10} L_{10}
$S_{1,0}$	$10^2/1 \cdot 5$	$2 \cdot 10/1^2$	K'_{11} L_{11}
$U_{1,0}$	$9^2/1 \cdot 3$	$3 \cdot 9/1^2$	M_{11}
$J'_{2,0}$	$2^2 10^2/1^2 5^2$	$2^2 10^2/1^2 5^2$	$J'_{2,0}$
$L_{1,0}$ $K'_{1,0}$	$2 \cdot 8^2/1^2 4$	$2 \cdot 8^2/1^2 4$	$L_{1,0}$ $K'_{1,0}$
$M_{1,0}$	$7^2/1^2$	$7^2/1^2$	$M_{1,0}$
$I_{1,0}$	$3^2 6^2/1^2 2^2$	$3^2 6^2/1^2 2^2$	$I_{1,0}$

Table 8: The duality: 8 triangle ICIS versus 8 bimodal series

Name	μ	Dol	Gab	d	h	h^*	μ^*	Dual
J'_9	9	2 3 10	2 2 2 3	-4	18	144	15	$J_{3,-1}$
J'_{10}	10	2 4 8	2 2 2 4	8	14	84	14	$Z_{1,-1}$
J'_{11}	11	3 3 7	2 2 2 5	-12	12	60	13	$Q_{2,-1}$
K'_{10}	10	2 6 6	2 3 2 3	12	12	60	14	$W_{1,-1}$
K'_{11}	11	3 5 5	2 3 2 4	-20	10	40	13	$S_{1,-1}$
L_{10}	10	2 5 7	2 2 3 3	11	12	132	14	$W_{1,-1}^\sharp$
L_{11}	11	3 4 6	2 2 3 4	-18	10	90	13	$S_{1,-1}^\sharp$
M_{11}	11	4 4 5	2 3 3 3	-24	9	72	13	$U_{1,-1}$

of rank 24. This lattice can be considered as the full homology lattice of a K3 surface,

$$K_{24} = H_0(S, \mathbb{Z}) \oplus H_2(S, \mathbb{Z}) \oplus H_4(S, \mathbb{Z}),$$

where the inner product on $H_0(S, \mathbb{Z}) \oplus H_4(S, \mathbb{Z})$ is defined in such a way that this lattice corresponds to a unimodular hyperbolic plane U . The orthogonal complement of H is the lattice $\tilde{H}$ of the singularity X^* (cf. Table 6).

Let us consider Pinkham's interpretation in the new cases. The Milnor fibre of a triangle or quadrilateral isolated hypersurface or complete intersection singularity can be compactified in such way that after resolving the singularities one gets a K3 surface S [P3]. We consider the dual graph of the curve configuration at infinity in each case. Let $G(p_1, p_2, p_3, p_4)$ and $\tilde{G}(p_1, p_2, p_3, p_4)$ be the subgraphs of the graphs of Fig. 4 and Fig. 2 respectively obtained by omitting the vertices $\mu^b - 1$ and μ^b , and $p_1 + p_2 - 1$, $p_1 + p_2 + 1$, and ν respectively. Denote by $M(p_1, p_2, p_3, p_4)$ and $\tilde{M}(p_1, p_2, p_3, p_4)$ the corresponding lattices. Recall that the homology lattice $H_2(S, \mathbb{Z})$ of the K3 surface is denoted by L .

First start with a triangle ICIS (X, x_0) with Dolgachev numbers (b_1, b_2, b_3) . Then the dual graph is the graph $G(b_1, b_2, b_3)$. This yields an embedding $M(b_1, b_2, b_3) \subset L$ and the orthogonal complement is the Milnor lattice $H = \tilde{M}(p_1, p_2, p_3, p_4) \oplus U$. By alternative (1) (cf. [EW]) the dual of (X, x_0) is a bimodal series; the Milnor lattice of the "virtual" $l = -1$ element of the corresponding series is $M(b_1, b_2, b_3) \oplus U$. One can even associate a Coxeter element to the dual "virtual" singularity; it has order $\text{lcm}(N_1, N_2)$ where N_1, N_2 are the degrees of the equations of (X, x_0) [EW], whereas the monodromy operator of (X, x_0) has order N_2 . There is no Saito duality of characteristic polynomials. The dual singularities and orders of the monodromy operators are listed in Table 8.

On the other hand, we can start with a quadrilateral hypersurface singularity (X, x_0) with Dolgachev numbers (b_1, b_2, b_3, b_4) . Then the dual graph is the graph $G(b_1, b_2, b_3, b_4)$. We obtain an embedding $M(b_1, b_2, b_3, b_4) \subset L$ and the orthogonal complement is the Milnor lattice of (X, x_0) described in Table 4. Here we use alternative (2) for the duality. The reduced Milnor lattice H^b of the dual triangle ICIS according to Table 4 is the lattice $M(b_1, b_2, b_3, b_4) \oplus U$.

Finally, let (X, x_0) be one of the 5 quadrilateral ICIS. Then the dual graph is again the graph $G(b_1, b_2, b_3, b_4)$. One has an embedding $M(b_1, b_2, b_3, b_4) \subset L$ and the orthogonal complement is the Milnor lattice of (X, x_0) described in Table 5. Combining both alternatives (1) and (2), the lattice $M(b_1, b_2, b_3, b_4) \oplus U$

Table 9: The duality between the quadrilateral ICIS

Name	μ	Dol	Gab	d	μ^*	Dual
$J'_{2,0}$	13	2 2 2 6	2 2 2 <u>6</u>	-16	11	$J'_{2,-1}$
$L_{1,0}$	13	2 2 3 5	2 2 3 <u>5</u> 2 2 <u>4</u> <u>4</u>	-28	11	$L_{1,-1}^\sharp$ $K_{1,-1}^\flat$
$K'_{1,0}$	13	2 2 4 4	2 3 2 <u>5</u> 2 <u>4</u> 2 <u>4</u>	-32	11	$L_{1,-1}$ $K'_{1,-1}$
$M_{1,0}$	13	2 3 3 4	2 3 <u>3</u> <u>4</u> 2 <u>3</u> <u>3</u> <u>4</u>	-42	11	$M_{1,-1}^\sharp$ $M_{1,-1}$
$I_{1,0}$	13	3 3 3 3	<u>3</u> <u>3</u> <u>3</u> <u>3</u>	-54	11	$I_{1,-1}$

can be interpreted as follows: The 5 quadrilateral ICIS are the initial $l = 0$ elements of 8 series of ICIS. To each such series one can again associate a virtual $l = -1$ element with a well-defined Milnor lattice. Then $M(b_1, b_2, b_3, b_4) \oplus U$ is the reduced Milnor lattice $\tilde{H}^b$ of the dual virtual singularity. This correspondence is indicated in Table 9. There is also a duality between the 8 virtual singularities as indicated in [EW] (see also [E3, Table 3.6.2]). There is no Saito duality of characteristic polynomials in both cases. But as we have seen above, using alternative (2) we get a third correspondence, for which Saito's duality of characteristic polynomials holds.

By Dolgachev's construction [D4], to each case of Pinkham's construction there corresponds a pair of mirror symmetric families of K3 surfaces. Moreover, also to each case where we only have a pair of lattices embedded as orthogonal complements to each other in the lattice K_{24} (cf. Table 6) there corresponds such a mirror pair.

One can also investigate Kobayashi's duality of weight systems for our extension of Arnold's strange duality. As already observed by Kobayashi [Kob], only some of the weight systems of the quadrilateral hypersurface singularities have dual weight systems, the dual weight systems are in general not unique and they correspond again to isolated hypersurface singularities. Since an ICIS has two degrees N_1 and N_2 , it is not quite clear how to generalize the notion of a weighted magic square. One possibility would be to work with the sum of the degrees $N := N_1 + N_2$ and to use also 3×4 and 4×4 matrices instead of 3×3 matrices. Then one finds again that in some cases there does not exist a dual weight system, the dual weight systems are in general not unique, most cases are self-dual, and only in the cases $J'_{10} \leftrightarrow Z_{1,0}$, $K'_{11} \leftrightarrow S_{1,0}$, $J'_{2,0} \leftrightarrow J'_{2,0}$, and $M_{1,0} \leftrightarrow M_{1,0}$ of our duality there exist weighted magic squares giving a duality of the corresponding weight systems.

However, there is a relation between our extended duality and a polar duality between the Newton polytopes generalizing Kobayashi's observation for Arnold's strange duality. This can be used to explain Saito's duality of characteristic polynomials. For details see the forthcoming paper [E6].

5 Singular moonshine

Let us consider the symbols $\pi\pi^*$ of Table 7. It turns out that they all occur in the list of Kondo, too. These pairs and the pairs from the original Arnold

Table 10: Self-dual Frame shapes of G with trace -2 , -3 or -4

ATLAS	Frame		Niemeier	Duality
21A	$2^2 3^2 7^2 42^2 / 1^2 6^2 14^2 21^2$	*		$E_{12} \leftrightarrow E_{12}$
15E	$2^2 3 \cdot 5 \cdot 30^2 / 1^2 6 \cdot 10 \cdot 15^2$	*	$D_{16} \oplus E_8$	$E_{13} \leftrightarrow Z_{11}$
24B	$2 \cdot 3^2 4 \cdot 24^2 / 1^2 6 \cdot 8^2 12$	*		$E_{14} \leftrightarrow Q_{10}$
11A	$2^2 22^2 / 1^2 11^2$	*	D_{12}^2	$Z_{12} \leftrightarrow Z_{12}$
18B	$2 \cdot 3 \cdot 18^2 / 1^2 6 \cdot 9$	*	$A_{17} \oplus E_7$	$Z_{13} \leftrightarrow Q_{11}$
15B	$3^2 15^2 / 1^2 5^2$	*		$Q_{12} \leftrightarrow Q_{12}$
20A	$2^2 5^2 20^2 / 1^2 4^2 10^2$	*		$W_{12} \leftrightarrow W_{12}$
16B	$2 \cdot 16^2 / 1^2 8$	*	$A_{15} \oplus D_9$	$W_{13} \leftrightarrow S_{11}$
13A	$13^2 / 1^2$	*	A_{12}^2	$S_{12} \leftrightarrow S_{12}$
12E	$4^2 12^2 / 1^2 3^2$	*		$U_{12} \leftrightarrow U_{12}$
9C	$2^3 3^2 18^3 / 1^3 6^2 9^3$		$D_{10} \oplus E_7^2$	$J'_9 \leftrightarrow J_{3,0}$
7B	$2^3 14^3 / 1^3 7^3$	*	D_8^3	$J'_{10} \leftrightarrow Z_{1,0}$
12K	$2^2 3 \cdot 12^3 / 1^3 4 \cdot 6^2$	*	$A_{11} \oplus D_7 \oplus E_6$	$J'_{11} \leftrightarrow Q_{2,0}$ $K'_{10} \leftrightarrow W_{1,0}$ $L_{10} \leftrightarrow W_{1,0}$
10E	$2 \cdot 10^3 / 1^3 5$	*	$A_9^2 \oplus D_6$	$K'_{11} \leftrightarrow S_{1,0}$ $L_{11} \leftrightarrow S_{1,0}$
9A	$9^3 / 1^3$		A_8^3	$M_{11} \leftrightarrow U_{1,0}$
5B	$2^4 10^4 / 1^4 5^4$		D_6^4	$J'_{2,0} \leftrightarrow J'_{2,0}$
8C	$2^2 8^4 / 1^4 4^2$		$A_7^2 \oplus D_5^2$	$L_{1,0} \leftrightarrow L_{1,0}$ $L_{1,0} \leftrightarrow K'_{1,0}$ $K'_{1,0} \leftrightarrow K'_{1,0}$
7A	$7^4 / 1^4$		A_6^4	$M_{1,0} \leftrightarrow M_{1,0}$
6A	$3^4 6^4 / 1^4 2^4$			$I_{1,0} \leftrightarrow I_{1,0}$
10A	$5^2 10^2 / 1^2 2^2$	*		
15A	$2^3 3^3 5^3 30^3 / 1^3 6^3 10^3 15^3$	*	E_8^3	
12A	$2^4 3^4 12^4 / 1^4 4^4 6^4$		E_6^4	

duality correspond to self-dual Frame shapes of the group G with trace -2 , -3 , or -4 . By examining Kondo's list one finds that there are 22 such Frame shapes and all but 3 occur. They are listed in Table 10. Here we use the ATLAS notation [ATL] for the conjugacy classes. For each value of the trace one symbol is missing.

Special elements of G correspond to the deep holes of the Leech lattice. A *deep hole* of the Leech lattice Λ_{24} is a point in $\mathbb{R}^{24}$ which has maximal distance from the lattice points. It is a beautiful theorem of J. H. Conway, R. A. Parker, and N. J. A. Sloane ([CPS], see also [E4]) that there are 23 types of deep holes in Λ_{24} which are in one-to-one correspondence with the 23 isomorphism classes of even unimodular lattices in $\mathbb{R}^{24}$ containing roots, which were classified by H.-V. Niemeier [Nie]. These lattices are characterized by the root systems which they contain. The Frame shape of the Coxeter element of such a root system is also the Frame shape of an automorphism of the Leech lattice. We have indicated in Table 10 the type of the root system of the Niemeier lattice, if a Frame shape corresponds to the Coxeter element of such a root system.

The automorphism group G of the Leech lattice contains the Mathieu group

M_{24} (see e.g. [E4]). S. Mukai [M] has classified the finite automorphism groups of K3 surfaces (automorphisms which leave the symplectic form invariant) and shown that they admit a certain embedding into the Mathieu group M_{24} . He gives a list of 11 maximal groups such that every finite automorphism group imbeds into one of these groups. A table of the centralizers of the conjugacy classes of G can be found in [Wi]. For an element $g \in G$, denote its centralizer by $Z(g)$ and the finite cyclic group generated by g by $\langle g \rangle$. We have marked by (*) in Table 10 the cases where there is an obvious inclusion of $Z(g)/\langle g \rangle$ in one of Mukai's groups. It follows that in these cases there is a K3 surface with an operation of $Z(g)/\langle g \rangle$ on it by symplectic automorphisms.

To a Frame shape

$$\pi = \prod_{m|N} m^{\chi_m}$$

one can associate a modular function [Kon]. Let

$$\eta(\tau) = q^{1/24} \prod_{n=1}^{\infty} (1 - q^n), \quad q = e^{2\pi i \tau}, \tau \in \mathbb{H},$$

be the Dedekind η -function. Then define

$$\eta_{\pi}(\tau) = \prod_{m|N} \eta(m\tau)^{\chi_m}.$$

Saito [S] has proved the identity

$$\eta_{\pi} \left(-\frac{1}{N\tau} \right) \eta_{\pi^*}(\tau) \sqrt{d} = 1,$$

where $d = \prod m^{\chi_m}$ and π^* is the dual Frame shape. From this it follows that $\eta_{\pi\pi^*}$ is a modular function for the group

$$\Gamma_0(N) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z}) \mid c \equiv 0(N) \right\}.$$

Question 1 Let $\pi\pi^*$ be one of the self-dual Frame shapes of Table 10. Is there any relation of $\eta_{\pi\pi^*}$ to the analogue of Dedekind's eta function for K3 surfaces considered in [JT] ?

The Frame shape $\pi\pi^*$ is the Frame shape of the operator $c \oplus c^*$ which can be considered as an automorphism of a sublattice of finite index of the even unimodular 24-dimensional lattice K_{24} , which is the full homology lattice of a K3 surface. The lattice K_{24} has the same rank as the Leech lattice, but contrary to the Leech lattice it is indefinite and has signature $(4, 20)$.

Question 2 Is there an explanation for this strange correspondence between operators of different lattices?

Is it only a purely combinatorial coincidence? One can try to classify finite sequences $(\chi_1, \chi_2, \dots, \chi_N)$ with the following properties:

- (1) $\chi_m \in \mathbb{Z}$ for all $m = 1, \dots, N$,

- (2) $\sum m\chi_m = 24$,
- (3) $\chi_m = 0$ for $m \nmid N$,
- (4) $\chi_m = -\chi_{N/m}$ for $m|N$,
- (5) $\prod m^{\chi_m} \in \mathbb{N}$,
- (6) $\chi_1 \in \{-2, -3, -4\}$,
- (7) $|\chi_m| \leq |\chi_N|$ for $m|N$.

By a computer search, one finds for $N \leq 119$ in addition to the 22 Frame shapes of Table 10 only the following Frame shapes:

$$3^2 6 \cdot 12^2 / 1^2 2 \cdot 4^2, \quad 2^4 3^4 4^4 24^4 / 1^4 6^4 8^4 12^4, \quad 2^2 4^2 5^2 40^2 / 1^2 8^2 10^2 20^2.$$

These Frame shapes also appear in Kondo's tables, namely in the tables of certain transforms of the Frame shapes of G [Kon, Table III, 4C; Table II, 12A; Table II, 20A].

References

- [AC1] N. A'Campo: Le nombre de Lefschetz d'une monodromie. *Indag. Math.* **35**, 113–118 (1973).
- [AC2] N. A'Campo: La fonction zêta d'une monodromie. *Comment. Math. Helvetici* **50**, 233–248 (1975).
- [Ar] V. I. Arnold: Critical points of smooth functions and their normal forms. *Usp. Math. Nauk.* **30**:5, 3–65 (1975) (Engl. translation in *Russ. Math. Surv.* **30**:5, 1–75 (1975)).
- [AGV] V. I. Arnold, S. M. Gusein-Zade, A. N. Varchenko: *Singularities of Differentiable Maps, Volume II*. Birkhäuser, Boston Basel Berlin 1988.
- [Ba] V. V. Batyrev: Dual polyhedra and the mirror symmetry for Calabi-Yau hypersurfaces in toric varieties. *J. Alg. Geom.* **3**, 493–535 (1994).
- [Br] E. Brieskorn: Die Monodromie der isolierten Singularitäten von Hyperflächen. *Manuscripta math.* **2**, 103–161 (1970).
- [ATL] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker, R. A. Wilson: *ATLAS of Finite Groups*. Oxford Univ. Press, London New York 1985.
- [CN] J. H. Conway, S. P. Norton: Monstrous moonshine. *Bull. Lond. Math. Soc.* **11**, 308–339 (1979).
- [CPS] J. H. Conway, R. A. Parker, N. J. A. Sloane: The covering radius of the Leech lattice. *Proc. R. Soc. Lond. A* **380**, 261–290 (1982).
- [D1] I. V. Dolgachev: Quotient-conical singularities on complex surfaces. *Funkt. Anal. Jego Prilozh.* **8**:2, 75–76 (1974) (Engl. translation in *Funct. Anal. Appl.* **8**, 160–161 (1974)).

- [D2] I. V. Dolgachev: Automorphic forms and weighted homogeneous singularities. *Funkt. Anal. Jego Prilozh.* **9**:2, 67–68 (1975) (Engl. translation in *Funct. Anal. Appl.* **9**, 149–151 (1975)).
- [D3] I. V. Dolgachev: Integral quadratic forms: applications to algebraic geometry. In: *Sem. Bourbaki*, 1982/83, Exposé n° 611, *Astérisque*, Vol. **105/106**, Soc. Math. France 1983, pp. 251–275.
- [D4] I. V. Dolgachev: Mirror symmetry for lattice polarized K3 surfaces. Preprint, University of Michigan, Ann Arbor, 1995 (alg-geom/9502005).
- [DN] I. V. Dolgachev, V. V. Nikulin: Exceptional singularities of V. I. Arnold and K3 surfaces. In: *Proc. USSR Topological Conference in Minsk*, 1977.
- [E1] W. Ebeling: Quadratische Formen und Monodromiegruppen von Singularitäten. *Math. Ann.* **255**, 463–498 (1981).
- [E2] W. Ebeling: Milnor lattices and geometric bases of some special singularities. *L'Enseignement Math.* **29**, 263–280 (1983).
- [E3] W. Ebeling: The Monodromy Groups of Isolated Singularities of Complete Intersections. *Lect. Notes in Math.*, Vol. **1293**, Springer-Verlag, Berlin etc., 1987.
- [E4] W. Ebeling: *Lattices and Codes*. Vieweg, Wiesbaden 1994.
- [E5] W. Ebeling: On Coxeter-Dynkin diagrams of hypersurface singularities. *J. Math. Sciences* **82** (5), 3657–3664 (1996).
- [E6] W. Ebeling: Strange duality and polar duality. In preparation.
- [EW] W. Ebeling, C. T. C. Wall: Kodaira singularities and an extension of Arnold's strange duality. *Compositio Math.* **56**, 3–77 (1985).
- [F] J. S. Frame: Computation of the characters of the Higman-Sims group and its automorphism group. *J. Algebra* **20** (2), 320–349 (1972).
- [G] A. M. Gabrielov: Dynkin diagrams of unimodal singularities. *Funkt. Anal. Jego Prilozh.* **8**:3, 1–6 (1974) (Engl. translation in *Funct. Anal. Appl.* **8**, 192–196 (1974)).
- [H1] H. Hamm: Lokale topologische Eigenschaften komplexer Räume. *Math. Ann.* **191**, 235–252 (1971).
- [H2] H. Hamm: Exotische Sphären als Umgebungsränder in speziellen komplexen Räumen. *Math. Ann.* **197**, 44–56 (1972).
- [JT] J. Jorgenson, A. Todorov: A conjectured analogue of Dedekind's eta function for K3 surfaces. *Math. Res. Letters* **2**, 359–376 (1995).
- [Kob] M. Kobayashi: Duality of weights, mirror symmetry and Arnold's strange duality. Preprint Tokyo Institute of Technology 1995 (alg-geom/9502004).
- [Kon] T. Kondo: The automorphism group of Leech lattice and elliptic modular functions. *J. Math. Soc. Japan* **37**, 337–361 (1985).

- [L1] E. Looijenga: The smoothing components of a triangle singularity. I. Proc. Symp. Pure Math. Vol. **40**, Part 2, 173–183 (1983).
- [L2] E. Looijenga: The smoothing components of a triangle singularity. II. Math. Ann. **269**, 357–387 (1984).
- [M] S. Mukai: Finite groups of automorphisms of K3 surfaces and the Mathieu group. Invent. math. **94**, 183–221 (1988).
- [Na1] I. Nakamura: Inoue-Hirzebruch surfaces and a duality of hyperbolic unimodular singularities I. Math. Ann. **252**, 221–235 (1980).
- [Na2] I. Nakamura: Duality of cusp singularities. In: Complex Analysis of Singularities (R.I.M.S. Symposium, K. Aomoto, organizer), Kyoto 1981, pp. 1–18.
- [Nie] H.-V. Niemeier: Definite quadratische Formen der Dimension 24 und Diskriminante 1. J. Number Theory **5**, 142–178 (1973).
- [Nik] V. V. Nikulin: Integral symmetric bilinear forms and some of their applications. Izv. Akad. Nauk. SSSR Ser. Mat. **43**, 111–177 (1979) (Engl. translation in Math. USSR Izv. **14**, No. 1, 103–167 (1980)).
- [OW] P. Orlik, P. Wagreich: Isolated singularities of algebraic surfaces with $\mathbb{C}^*$ -action. Ann. of Math. **93**, 205–228 (1971).
- [P1] H. Pinkham: Singularités exceptionnelles, la dualité étrange d’Arnold et les surfaces K-3. C. R. Acad. Sc. Paris, Serie A, **284**, 615–618 (1977).
- [P2] H. Pinkham: Groupe de monodromie des singularités unimodulaires exceptionnelles. C. R. Acad. Sc. Paris, Serie A, **284**, 1515–1518 (1977).
- [P3] H. Pinkham: Deformations of normal surface singularities with $\mathbb{C}^*$ action. Math. Ann. **232**, 65–84 (1978).
- [S] K. Saito: On a duality of characteristic polynomials for regular systems of weights. Preprint RIMS, Kyoto University 1994.
- [Wa] C. T. C. Wall: Classification of unimodal isolated singularities of complete intersections. Proc. Symp. Pure Math. Vol. **40**, Part 2, 625–640 (1983).
- [Wi] R. A. Wilson: The maximal subgroups of Conway’s group Co_1 . J. Algebra **85**, 144–165 (1983).

INSTITUT FÜR MATHEMATIK, UNIVERSITÄT HANNOVER, POSTFACH 6009,
D-30060 HANNOVER, GERMANY
E-mail address: ebeling@math.uni-hannover.de